



SAMPLE REPORT

Case Description: Karen Z. — Inpatient Mental Health Interpretive Report

Karen Z., a 35-year-old married woman, mother of two children, was admitted to a private inpatient psychiatric unit after her second suicide attempt. Although she has been taking antidepressant medication, her depressed moods have persisted. Two months earlier, her psychiatrist prescribed amitriptyline, which she reported did not help her mood and was extremely sedating.

Ms Z. has a history of mental health contacts since she was 18 years of age, when she first sought help for her mood swings. At that time she was treated with antidepressants. Her depression became more severe three years ago, when she made her first suicide attempt. At that time, she underwent electroconvulsive therapy (ECT) after her medication failed to provide relief for her depression. She was able to return to work after her treatment.

Ms Z. completed high school and a two-year associate's degree program at a community college. She has been employed in office management for a large insurance firm. Her husband, who is employed by the same company, is very supportive of her and assumes most of the home and family responsibilities during her depression.

Karen Z., a 35-year-old married woman, mother of two children, was admitted to a private inpatient psychiatric unit after her second suicide attempt. Although she has been taking antidepressant medication, her depressed moods have persisted. Two months earlier, her psychiatrist prescribed amitriptyline, which she reported did not help her mood and was extremely sedating.

Ms Z. has a history of mental health contacts since she was 18 years of age, when she first sought help for her mood swings. At that time she was treated with antidepressants. Her depression became more severe three years ago, when she made her first suicide attempt. At that time, she underwent electroconvulsive therapy (ECT) after her medication failed to provide relief for her depression. She was able to return to work after her treatment.

Case descriptions do not accompany MMPI-2 reports, but are provided here as background information. The following report was generated from Q-global™, Pearson's web-based scoring and reporting application, using Ms. Z.'s responses to the MMPI-2. Additional MMPI-2 sample reports, product offerings, training opportunities, and resources can be found at PearsonClinical.com/mmpi2.

Copyright © 2014 Pearson Education, Inc. or its affiliate(s). All rights reserved. Q-global, Always Learning, Pearson, design for Psi, and PsychCorp are trademarks, in the U.S. and/or other countries, of Pearson Education, Inc. or its affiliate(s). Minnesota Multiphasic Personality Inventory-2 and MMPI-2 are registered trademarks of the University of Minnesota, Minneapolis, MN. 8795-A 01/14



SAMPLE REPORT

Case Description (*continued*): Karen Z. — Inpatient Mental Health Interpretive Report

Ms Z. completed high school and a two-year associate's degree program at a community college. She has been employed in office management for a large insurance firm. Her husband, who is employed by the same company, is very supportive of her and assumes most of the home and family responsibilities during her depression.



Inpatient Mental Health Interpretive Report

MMPI®-2

The Minnesota Report™: Adult Clinical System-Revised, 4th Edition

James N. Butcher, PhD

Name:	Karen Z.
ID Number:	2513
Age:	35
Gender:	Female
Marital Status:	Married
Years of Education:	14
Date Assessed:	1/31/14



Copyright © 1989, 1993, 2001, 2005 by the Regents of the University of Minnesota. All rights reserved.
Portions reproduced from the MMPI-2 test booklet. Copyright © 1942, 1943 (renewed 1970), 1989 by the Regents of the University of Minnesota. All rights reserved. Portions excerpted from the *MMPI-2 Manual for Administration, Scoring, and Interpretation, Revised Edition*. Copyright © 2001 by the Regents of the University of Minnesota. All rights reserved.
Distributed exclusively under license from the University of Minnesota by NCS Pearson, Inc.


Minnesota Multiphasic Personality Inventory and **MMPI** are registered trademarks and **The Minnesota Report** is a trademark of the University of Minnesota. **Pearson**, the **PSI logo**, and **PsychCorp** are trademarks in the U.S. and/or other countries of Pearson Education, Inc., or its affiliate(s).

TRADE SECRET INFORMATION

Not for release under HIPAA or other data disclosure laws that exempt trade secrets from disclosure.

[9.5 / 1 / QG]

MMPI-2 VALIDITY PATTERN




Raw Score:	6	7	14	9	2	2	17	31
T Score:	54	65 ^F	85	77	57	42	54	57
Response %:	100	100	100	100	100	100	100	100

Cannot Say (Raw): 0
 Percent True: 37
 Percent False: 63

	Raw Score	T Score	Resp. %
S ₁ - Beliefs in Human Goodness	15	69	100
S ₂ - Serenity	3	41	100
S ₃ - Contentment with Life	4	52	100
S ₄ - Patience/Denial of Irritability	4	48	100
S ₅ - Denial of Moral Flaws	5	61	100

MMPI-2 CLINICAL AND SUPPLEMENTARY SCALES PROFILE



Raw Score:	21	49	38	22	34	15	30	35	12	50	15	24	1	26	7	7
K Correction:	9			7			17	17	3							
T Score:	86	112	87	66	55	67	86	90	41	74	40	52	44	77	37	68
Response %:	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100

Welsh Code: 2**8*317"0'64+-5/9: F""+-K/L:

Profile Elevation: 79.4

PROFILE VALIDITY

This MMPI-2 profile should be interpreted with caution. There is some possibility that the clinical report is an exaggerated picture of the client's present situation and problems. She is presenting an unusual number of psychological symptoms. This response set could result from poor reading ability, confusion, disorientation, stress, or a need to seek a great deal of attention for her problems.

SYMPTOMATIC PATTERNS

This report was developed using the D and Sc scales as the prototype. The client is quite depressed at this time. In addition, she exhibits a chronic pattern of alienation and isolation. She appears to be quite tense, anxious, and withdrawn. She is having a great deal of difficulty concentrating and cannot keep her mind on tasks. She appears to worry a lot and is functioning at a low level of efficiency as a result of her depressed mood. She may seem lethargic and apathetic and may show little interest in life. Life seems to have little meaning for her and she has scarce energy left over from mere survival to expend on any pleasures. Individuals with this profile often appear to be unable to feel pleasure as most people can. Life is drudgery for them.

Her emotional life is flat and blunted. She may show some signs of a thought disorder in addition to an affective disorder. Many individuals with this profile consider committing suicide and she may actually have serious plans for self-destruction. Individuals with this MMPI-2 clinical profile tend to be faced with multiple problems resulting from a long-standing lack of achievement and ineffective work behavior.

In addition, the following description is suggested by the client's scores on the content scales. She endorsed a number of items suggesting that she is experiencing low morale and a depressed mood. She reports a preoccupation with feeling guilty and unworthy. She feels that she deserves to be punished for wrongs she has committed. She feels regretful and unhappy about life, and she seems plagued by anxiety and worry about the future. She feels hopeless at times and feels that she is a condemned person. She endorsed response content that reflects low self-esteem and long-standing beliefs about her inadequacy. She has difficulty managing routine affairs, and the items she endorsed suggest a poor memory, concentration problems, and an inability to make decisions. She appears to be immobilized and withdrawn and has no energy for life. She views her physical health as failing and reports numerous somatic concerns. She feels that life is no longer worthwhile and that she is losing control of her thought processes.

According to her response content, there is a strong possibility that she has seriously contemplated suicide. She acknowledged having suicidal thoughts recently. Although she denies suicidal attempts in the past, given her current mood, an evaluation of suicidal potential appears to be indicated. She is rather high-strung and believes that she feels things more, or more intensely, than others do. She feels quite lonely and misunderstood at times.

Long-term personality factors identified by her PSY-5 scale elevations may help provide a clinical context for the symptoms she is presently experiencing. She shows a meager capacity to experience pleasure in life. Persons with high scores on INTR (Introversion/Low Positive Emotionality) tend to be pessimistic.

PROFILE FREQUENCY

Profile interpretation can be greatly facilitated by examining the relative frequency of clinical scale patterns in various settings. The client's high-point clinical scale score (D) occurred in 7.0% of the MMPI-2 normative sample of women. However, only 4.4% of the women had D scale peak scores at or above a T score of 65, and only 2.1% had well-defined D spikes. This high MMPI-2 profile configuration (2-8/8-2) is very rare in samples of normals, occurring in less than 1% of the MMPI-2 normative sample of women.

This high-point MMPI-2 score is very frequent among various samples of inpatient women. In the Graham and Butcher (1988) sample, the D scale occurred as the high point in 12.2% of the females (all of the cases were at or above a T score of 65, and 8.4% were well defined in that range). In the Pearson inpatient sample, the high-point clinical scale score on D occurred in 25.4% of the women. Moreover, 24.3% had the D scale spike at or over a T score of 65, and 13.8% produced well-defined D scale peak scores in that range.

Her high MMPI-2 profile configuration (2-8/8-2) was found in 4.6% of the females in the Graham and Butcher (1988) sample and in 4.0% of the females in the Pearson inpatient sample.

PROFILE STABILITY

The relative elevation of her highest clinical scale scores suggests some lack of clarity in profile definition. Although her most elevated clinical scales are likely to be present in her profile pattern if she is retested at a later date, there could be some shifting of the most prominent scale elevations in the profile code. The difference between the profile type used to develop the present report (involving D and Sc) and the next highest scale in the profile code was 3 points. So, for example, if the client is tested at a later date, her profile might involve more behavioral elements related to elevations on Hy. If so, then on retesting, intensification of defense mechanisms such as repression and denial might occur along with the presentation of vague physical problems.

INTERPERSONAL RELATIONS

She has great problems with alienation in social relationships. She feels vulnerable to interpersonal hurt, lacks trust, and may never form close, satisfying interpersonal ties. She feels very insecure in relationships and may be preoccupied with guilt and self-defeating behavior. Her marital situation is likely to be unrewarding and impoverished. Married individuals with this profile typically are insecure in relationships. They lack confidence in their marriage and feel that something bad is going to happen to them. The client is likely to feel that things have gone wrong in her marriage.

She is a very introverted person who has difficulty meeting and interacting with other people. She is shy and emotionally distant. She tends to be very uneasy, rigid, and overcontrolled in social situations. Her shyness is probably symptomatic of a broader pattern of social withdrawal. Personality characteristics related to social introversion tend to be stable over time. Her generally reclusive behavior, introverted lifestyle, and tendency toward interpersonal avoidance may be prominent in any future test results.

Her high score on the Marital Distress Scale suggests that her marital situation is problematic at this time. She reported a number of problems with her marriage that are possibly important to understanding her current psychological symptoms.

The client's scores on the content scales suggest the following additional information concerning her interpersonal relations. Her social relationships are likely to be viewed by others as problematic. She may be visibly uneasy around others, sits alone in group situations, and dislikes engaging in group activities.

DIAGNOSTIC CONSIDERATIONS

Similar clients tend to have features of both an affective disorder and a thought disorder. In addition, there seems to be a long-standing pattern of maladjustment that is characteristic of people with severe personality disorders. She is likely to be diagnosed as having a Schizophrenic or Major Affective Disorder. Her scores on the content scales indicate that her acknowledged tendency toward experiencing depressed mood should be taken into consideration in any diagnostic formulation.

TREATMENT CONSIDERATIONS

Multiple problems and difficulties forming interpersonal relationships make patients with this profile poor candidates for relationship-based psychotherapy. Their basic lack of trust and withdrawal also make psychotherapy difficult. Some outpatients with this profile seem to benefit from minimal contact treatment, such as brief periodic visits with a directive, supportive therapist. Many need psychotropic medication to control their bizarre thinking or to elevate their mood.

The client's scores on the content scales seem to indicate low potential for change. She may feel that her problems are not addressable through therapy and that she is not likely to benefit much from psychological treatment at this time. Her apparently negative treatment attitudes may need to be explored early in therapy if treatment is to be successful.

In any intervention or psychological evaluation program involving occupational adjustment, her negative work attitudes could become an important problem to overcome. She has a number of attitudes and feelings that could interfere with work adjustment.

ADDITIONAL SCALES

	Raw Score	T Score	Resp %
Personality Psychopathology Five (PSY-5) Scales			
Aggressiveness (AGGR)	0	30	100
Psychoticism (PSYC)	3	50	100
Disconstraint (DISC)	6	39	100
Negative Emotionality/Neuroticism (NEGE)	17	59	100
Introversion/Low Positive Emotionality (INTR)	30	99	100
Supplementary Scales			
Anxiety (A)	27	69	100
Repression (R)	29	83	100
Ego Strength (Es)	19	30	100
Dominance (Do)	13	39	100
Social Responsibility (Re)	24	59	100
Harris-Lingoes Subscales			
Depression Subscales			
Subjective Depression (D ₁)	29	101	100
Psychomotor Retardation (D ₂)	11	79	100
Physical Malfunctioning (D ₃)	9	93	100
Mental Dullness (D ₄)	14	102	100
Brooding (D ₅)	9	83	100
Hysteria Subscales			
Denial of Social Anxiety (Hy ₁)	1	35	100
Need for Affection (Hy ₂)	10	63	100
Lassitude-Malaise (Hy ₃)	14	95	100
Somatic Complaints (Hy ₄)	7	65	100
Inhibition of Aggression (Hy ₅)	4	54	100
Psychopathic Deviate Subscales			
Familial Discord (Pd ₁)	3	56	100
Authority Problems (Pd ₂)	3	53	100
Social Imperturbability (Pd ₃)	1	35	100
Social Alienation (Pd ₄)	6	60	100
Self-Alienation (Pd ₅)	6	63	100
Paranoia Subscales			
Persecutory Ideas (Pa ₁)	2	51	100
Poignancy (Pa ₂)	5	65	100
Naivete (Pa ₃)	8	65	100

	Raw Score	T Score	Resp %
Schizophrenia Subscales			
Social Alienation (Sc ₁)	8	69	100
Emotional Alienation (Sc ₂)	5	86	100
Lack of Ego Mastery, Cognitive (Sc ₃)	6	80	100
Lack of Ego Mastery, Conative (Sc ₄)	10	90	100
Lack of Ego Mastery, Defective Inhibition (Sc ₅)	2	53	100
Bizarre Sensory Experiences (Sc ₆)	9	81	100
Hypomania Subscales			
Amorality (Ma ₁)	2	54	100
Psychomotor Acceleration (Ma ₂)	5	50	100
Imperturbability (Ma ₃)	1	37	100
Ego Inflation (Ma ₄)	1	37	100
Social Introversion Subscales (Ben-Porath, Hostetler, Butcher, & Graham)			
Shyness/Self-Consciousness (Si ₁)	12	68	100
Social Avoidance (Si ₂)	8	74	100
Alienation--Self and Others (Si ₃)	7	55	100
Content Component Scales (Ben-Porath & Sherwood)			
Fears Subscales			
Generalized Fearfulness (FRS ₁)	1	48	100
Multiple Fears (FRS ₂)	2	37	100
Depression Subscales			
Lack of Drive (DEP ₁)	11	95	100
Dysphoria (DEP ₂)	6	79	100
Self-Depreciation (DEP ₃)	3	61	100
Suicidal Ideation (DEP ₄)	2	77	100
Health Concerns Subscales			
Gastrointestinal Symptoms (HEA ₁)	3	75	100
Neurological Symptoms (HEA ₂)	8	83	100
General Health Concerns (HEA ₃)	6	87	100
Bizarre Mentation Subscales			
Psychotic Symptomatology (BIZ ₁)	0	44	100
Schizotypal Characteristics (BIZ ₂)	0	41	100
Anger Subscales			
Explosive Behavior (ANG ₁)	2	54	100
Irritability (ANG ₂)	3	49	100
Cynicism Subscales			
Misanthropic Beliefs (CYN ₁)	0	34	100
Interpersonal Suspiciousness (CYN ₂)	1	40	100

	Raw Score	T Score	Resp %
Antisocial Practices Subscales			
Antisocial Attitudes (ASP ₁)	1	36	100
Antisocial Behavior (ASP ₂)	1	51	100
Type A Subscales			
Impatience (TPA ₁)	4	58	100
Competitive Drive (TPA ₂)	1	40	100
Low Self-Esteem Subscales			
Self-Doubt (LSE ₁)	7	71	100
Submissiveness (LSE ₂)	5	69	100
Social Discomfort Subscales			
Introversion (SOD ₁)	16	85	100
Shyness (SOD ₂)	6	65	100
Family Problems Subscales			
Family Discord (FAM ₁)	2	42	100
Familial Alienation (FAM ₂)	0	41	100
Negative Treatment Indicators Subscales			
Low Motivation (TRT ₁)	9	86	100
Inability to Disclose (TRT ₂)	3	60	100

Uniform T scores are used for Hs, D, Hy, Pd, Pa, Pt, Sc, Ma, the content scales, the content component scales, and the PSY-5 scales. The remaining scales and subscales use linear T scores.

CRITICAL ITEMS

The following critical items have been found to have possible significance in analyzing a client's problem situation. Although these items may serve as a source of hypotheses for further investigation, caution should be used in interpreting individual items because they may have been checked inadvertently.

The percentages of endorsement for each critical item by various reference groups are presented in brackets following the listing of the item. The endorsement percentage labeled "N" is the percentage of the MMPI-2 normative sample of 1,462 women who endorsed the item in the scored direction. Endorsement percentages for the normative sample are reported for all critical items. When available, endorsement percentages for the setting are also reported. The designation "Ip1" refers to a sample of 131 female inpatients (Graham & Butcher, 1988); "Ip2" refers to a sample of 5,942 female inpatients (Pearson, 1993).

Acute Anxiety State (Koss-Butcher Critical Items)

Of the 17 possible items in this section, 12 were endorsed in the scored direction:

- 2. Item Content Omitted. (False) [N = 4; Ip1 = 30; Ip2 = 29]
- 3. Item Content Omitted. (False) [N = 34; Ip1 = 56; Ip2 = 68]
- 10. Item Content Omitted. (False) [N = 14; Ip1 = 47; Ip2 = 55]
- 15. Item Content Omitted. (True) [N = 36; Ip1 = 50; Ip2 = 56]
- 28. Item Content Omitted. (True) [N = 9; Ip1 = 28; Ip2 = 36]
- 59. Item Content Omitted. (True) [N = 11; Ip1 = 34; Ip2 = 38]
- 140. Item Content Omitted. (False) [N = 27; Ip1 = 53; Ip2 = 65]
- 218. Item Content Omitted. (True) [N = 24; Ip1 = 62; Ip2 = 58]
- 223. Item Content Omitted. (False) [N = 18; Ip1 = 51; Ip2 = 55]
- 301. Item Content Omitted. (True) [N = 20; Ip1 = 51; Ip2 = 56]
- 444. Item Content Omitted. (True) [N = 28; Ip1 = 56; Ip2 = 50]
- 469. Item Content Omitted. (True) [N = 31; Ip1 = 67; Ip2 = 75]



Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

Depressed Suicidal Ideation (Koss-Butcher Critical Items)

Of the 22 possible items in this section, 17 were endorsed in the scored direction:

- 9. Item Content Omitted. (False) [N = 18; Ip1 = 46; Ip2 = 57]
- 38. Item Content Omitted. (True) [N = 28; Ip1 = 64; Ip2 = 67]
- 65. Item Content Omitted. (True) [N = 9; Ip1 = 45; Ip2 = 53]
- 71. Item Content Omitted. (True) [N = 32; Ip1 = 70; Ip2 = 59]
- 75. Item Content Omitted. (False) [N = 5; Ip1 = 31; Ip2 = 28]
- 92. Item Content Omitted. (True) [N = 3; Ip1 = 27; Ip2 = 26]
- 95. Item Content Omitted. (False) [N = 11; Ip1 = 51; Ip2 = 58]
- 130. Item Content Omitted. (True) [N = 38; Ip1 = 70; Ip2 = 74]
- 146. Item Content Omitted. (True) [N = 46; Ip1 = 59; Ip2 = 65]
- 215. Item Content Omitted. (True) [N = 18; Ip1 = 43; Ip2 = 43]
- 233. Item Content Omitted. (True) [N = 32; Ip1 = 58; Ip2 = 56]
- 273. Item Content Omitted. (True) [N = 18; Ip1 = 60; Ip2 = 62]
- 388. Item Content Omitted. (False) [N = 36; Ip1 = 65; Ip2 = 73]
- 411. Item Content Omitted. (True) [N = 25; Ip1 = 62; Ip2 = 62]
- 454. Item Content Omitted. (True) [N = 4; Ip1 = 35; Ip2 = 32]
- 485. Item Content Omitted. (True) [N = 23; Ip1 = 61; Ip2 = 58]
- 506. Item Content Omitted. (True) [N = 4; Ip1 = 49; Ip2 = 40]



Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

Threatened Assault (Koss-Butcher Critical Items)

Of the 5 possible items in this section, 1 was endorsed in the scored direction:

- 37. Item Content Omitted. (True) [N = 38; Ip1 = 52; Ip2 = 53]

Situational Stress Due to Alcoholism (Koss-Butcher Critical Items)

Of the 7 possible items in this section, 2 were endorsed in the scored direction:

- 125. Item Content Omitted. (False) [N = 13; Ip1 = 56; Ip2 = 51]
- 502. Item Content Omitted. (True) [N = 21; Ip1 = 49; Ip2 = 43]



Special Note:
The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

Mental Confusion (Koss-Butcher Critical Items)

Of the 11 possible items in this section, 4 were endorsed in the scored direction:

- 31. Item Content Omitted. (True) [N = 13; Ip1 = 51; Ip2 = 55]
- 180. Item Content Omitted. (True) [N = 3; Ip1 = 41; Ip2 = 34]
- 299. Item Content Omitted. (True) [N = 15; Ip1 = 48; Ip2 = 47]
- 325. Item Content Omitted. (True) [N = 17; Ip1 = 53; Ip2 = 53]

Antisocial Attitude (Lachar-Wrobel Critical Items)

Of the 9 possible items in this section, 1 was endorsed in the scored direction:

- 35. Item Content Omitted. (True) [N = 37; Ip1 = 58; Ip2 = 42]

Family Conflict (Lachar-Wrobel Critical Items)

Of the 4 possible items in this section, 2 were endorsed in the scored direction:

- 21. Item Content Omitted. (True) [N = 41; Ip1 = 73; Ip2 = 65]
- 125. Item Content Omitted. (False) [N = 13; Ip1 = 56; Ip2 = 51]

Somatic Symptoms (Lachar-Wrobel Critical Items)

Of the 23 possible items in this section, 16 were endorsed in the scored direction:

- 28. Item Content Omitted. (True) [N = 9; Ip1 = 28; Ip2 = 36]
- 33. Item Content Omitted. (False) [N = 36; Ip1 = 53; Ip2 = 42]
- 40. Item Content Omitted. (True) [N = 7; Ip1 = 26; Ip2 = 32]

- 53. Item Content Omitted. (True) [N = 25; Ip1 = 52; Ip2 = 48]
- 57. Item Content Omitted. (False) [N = 32; Ip1 = 31; Ip2 = 54]
- 59. Item Content Omitted. (True) [N = 11; Ip1 = 34; Ip2 = 38]
- 101. Item Content Omitted. (True) [N = 7; Ip1 = 26; Ip2 = 29]
- 111. Item Content Omitted. (True) [N = 9; Ip1 = 25; Ip2 = 29]
- 142. Item Content Omitted. (False) [N = 7; Ip1 = 30; Ip2 = 22]
- 175. Item Content Omitted. (True) [N = 5; Ip1 = 36; Ip2 = 37]
- 176. Item Content Omitted. (False) [N = 26; Ip1 = 39; Ip2 = 51]
- 224. Item Content Omitted. (False) [N = 19; Ip1 = 35; Ip2 = 49]
- 229. Item Content Omitted. (True) [N = 5; Ip1 = 39; Ip2 = 27]
- 247. Item Content Omitted. (True) [N = 9; Ip1 = 25; Ip2 = 22]
- 295. Item Content Omitted. (False) [N = 16; Ip1 = 40; Ip2 = 31]
- 464. Item Content Omitted. (True) [N = 34; Ip1 = 56; Ip2 = 69]



Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

Sexual Concern and Deviation (Lachar-Wrobel Critical Items)

Of the 6 possible items in this section, 1 was endorsed in the scored direction:

- 12. Item Content Omitted. (False) [N = 26; Ip1 = 49; Ip2 = 52]

Anxiety and Tension (Lachar-Wrobel Critical Items)

Of the 11 possible items in this section, 7 were endorsed in the scored direction:

- 15. Item Content Omitted. (True) [N = 36; Ip1 = 50; Ip2 = 56]
- 218. Item Content Omitted. (True) [N = 24; Ip1 = 62; Ip2 = 58]
- 223. Item Content Omitted. (False) [N = 18; Ip1 = 51; Ip2 = 55]

- 261. Item Content Omitted. (False) [N = 48]
- 299. Item Content Omitted. (True) [N = 15; Ip1 = 48; Ip2 = 47]
- 301. Item Content Omitted. (True) [N = 20; Ip1 = 51; Ip2 = 56]
- 405. Item Content Omitted. (False) [N = 23]



Special Note:

The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

Sleep Disturbance (Lachar-Wrobel Critical Items)

Of the 6 possible items in this section, 4 were endorsed in the scored direction:

- 30. Item Content Omitted. (True) [N = 6]
- 140. Item Content Omitted. (False) [N = 27; Ip1 = 53; Ip2 = 65]
- 328. Item Content Omitted. (True) [N = 30]
- 471. Item Content Omitted. (True) [N = 18]

Deviant Thinking and Experience (Lachar-Wrobel Critical Items)

Of the 10 possible items in this section, 1 was endorsed in the scored direction:

- 122. Item Content Omitted. (True) [N = 82]

Depression and Worry (Lachar-Wrobel Critical Items)

Of the 16 possible items in this section, 14 were endorsed in the scored direction:

- 2. Item Content Omitted. (False) [N = 4; Ip1 = 30; Ip2 = 29]
- 3. Item Content Omitted. (False) [N = 34; Ip1 = 56; Ip2 = 68]
- 10. Item Content Omitted. (False) [N = 14; Ip1 = 47; Ip2 = 55]
- 65. Item Content Omitted. (True) [N = 9; Ip1 = 45; Ip2 = 53]
- 73. Item Content Omitted. (True) [N = 28]
- 75. Item Content Omitted. (False) [N = 5; Ip1 = 31; Ip2 = 28]
- 130. Item Content Omitted. (True) [N = 38; Ip1 = 70; Ip2 = 74]

165. Item Content Omitted. (False) [N = 10]
180. Item Content Omitted. (True) [N = 3; Ip1 = 41; Ip2 = 34]
273. Item Content Omitted. (True) [N = 18; Ip1 = 60; Ip2 = 62]
339. Item Content Omitted. (True) [N = 43]
411. Item Content Omitted. (True) [N = 25; Ip1 = 62; Ip2 = 62]
415. Item Content Omitted. (True) [N = 29]
454. Item Content Omitted. (True) [N = 4; Ip1 = 35; Ip2 = 32]



Special Note:
The content of the test items is included in the actual reports. To protect the integrity of the test, the item content does not appear in this sample report.

Deviant Beliefs (Lachar-Wrobel Critical Items)

Of the 15 possible items in this section, 1 was endorsed in the scored direction:

106. Item Content Omitted. (False) [N = 15]

Substance Abuse (Lachar-Wrobel Critical Items)

Of the 3 possible items in this section, 1 was endorsed in the scored direction:

168. Item Content Omitted. (True) [N = 8]

End of Report

NOTE: This MMPI-2 interpretation can serve as a useful source of hypotheses about clients. This report is based on objectively derived scale indices and scale interpretations that have been developed in diverse groups of patients. The personality descriptions, inferences, and recommendations contained herein need to be verified by other sources of clinical information because individual clients may not fully match the prototype. The information in this report should only be used by a trained and qualified test interpreter. The report was not designed or intended to be provided directly to clients. The information contained in the report is technical and was developed to aid professional interpretation.

This and previous pages of this report contain trade secrets and are not to be released in response to requests under HIPAA (or any other data disclosure law that exempts trade secret information from release). Further, release in response to litigation discovery demands should be made only in accordance with your profession's ethical guidelines and under an appropriate protective order.
